

On the index of a vector field tangent to a hypersurface with non-isolated zero in the embedding space

Oliver Klehn

Institut für Mathematik, Universität Hannover,
Postfach 6009, D-30060 Hannover, Germany
E-mail: klehn@math.uni-hannover.de

Abstract

We give a generalization of an algebraic formula of Gomez-Mont for the index of a vector field with isolated zero in $(\mathbb{C}^n, 0)$ and tangent to an isolated hypersurface singularity. We only assume that the vector field has an isolated zero on the singularity here.

1 Introduction

Let $(V, 0) = (\{f = 0\}, 0) \subseteq (\mathbb{C}^n, 0)$ be an isolated singularity of a hypersurface and

$$X = X_1 \frac{\partial}{\partial z_1} + \cdots + X_n \frac{\partial}{\partial z_n}$$

the germ of a holomorphic vector field on $(\mathbb{C}^n, 0)$ tangent to $(V, 0)$, say $Xf = cf$, with an isolated zero on $(V, 0)$. In [3] is defined a notion of an index $\text{ind}_{V,0} X$ of this vector field, called GSV index and generalizing the Poincaré-Hopf index. By contracting germs of Kähler forms on $(V, 0)$ with X one obtains two complexes

$$(\Omega_{V,0}^*, X): 0 \rightarrow \Omega_{V,0}^n \xrightarrow{X} \cdots \xrightarrow{X} \Omega_{V,0}^1 \xrightarrow{X} \mathcal{O}_{V,0} \rightarrow 0$$

$$(\Omega_{V,0}, X): 0 \rightarrow \Omega_{V,0}^{n-1} \xrightarrow{X} \cdots \xrightarrow{X} \Omega_{V,0}^1 \xrightarrow{X} \mathcal{O}_{V,0} \rightarrow 0$$

which have both finite dimensional homology spaces, see [2]. We set

$$h_i^* := \dim_{\mathbb{C}} H_i(\Omega_{V,0}^*, X) \quad h_i := \dim_{\mathbb{C}} H_i(\Omega_{V,0}, X).$$

It has been proved in [2] that the topologically defined GSV index is given as

$$\text{ind}_{V,0} X = \chi(\Omega_{V,0}, X) = \sum_{i=0}^{n-1} (-1)^i h_i.$$

Further the numbers h_i and h_i^* are computed under the condition that X has an isolated zero on $(\mathbb{C}^n, 0)$. We want to extend these computations to the case that X has an isolated zero only on $(V, 0)$ using the same methods as in [2].

Note that one can assume after a linear change of coordinates in $(\mathbb{C}^n, 0)$ that $(X_1, \dots, X_{n-1}, f)$ is a regular $\mathcal{O}$ -sequence. Throughout this paper we always assume the coordinates to be chosen in this way. We denote by

$$\mathcal{A} := \frac{\mathcal{O}}{(\partial f / \partial z_1, \dots, \partial f / \partial z_n)}$$

the Milnor algebra of f at 0 and we set

$$\mathcal{B} := \frac{\mathcal{O}}{(X_1, \dots, X_n)} \quad \mathcal{B}' := \frac{\mathcal{O}}{(X_1, \dots, X_{n-1})}.$$

We simply write $\mathcal{O}$ resp. Ω^i instead of $\mathcal{O}_{\mathbb{C}^n, 0}$ resp. $\Omega_{\mathbb{C}^n, 0}^i$. $\mathcal{B}'$ is infinite dimensional and $\mathcal{B}$ not necessarily finite dimensional. Our main result is the following:

Theorem 1. *The $\mathbb{C}$ -dimensions of the homology spaces of $(\Omega_{V,0}, X)$ are given by*

$$\begin{aligned} h_0^* &= \dim_{\mathbb{C}} \frac{\mathcal{B}}{(f)}, \\ h_1^* &= \dim_{\mathbb{C}} \frac{\text{ann}_{\mathcal{B}}(f)}{(c)} + \dim_{\mathbb{C}} \frac{\text{ann}_{\mathcal{B}'}(X_n)}{\text{ann}_{\mathcal{B}'}(X_n) \cap \mathcal{B}'(f, \partial f / \partial z_n)}, \\ h_2^* &= \dots = h_n^* = \dim_{\mathbb{C}} \frac{\text{ann}_{\mathcal{A}}(f)}{(c)} = \dim_{\mathbb{C}} \frac{\text{ann}_{\mathcal{A}}(c)}{(f)} =: \lambda. \end{aligned}$$

2 The homological index

The first steps are easy generalizations of the computations in [2]. Consider the complex

$$(\Omega^*, X): 0 \rightarrow \Omega^n \xrightarrow{X} \dots \xrightarrow{X} \Omega^1 \xrightarrow{X} \mathcal{O} \rightarrow 0.$$

Lemma 1. $H_j(\Omega^*, X) = 0$ for $j \geq 2$
 $H_1(\Omega^*, X) \cong \text{ann}_{\mathcal{B}'}(X_n)$, $(h_1 dz_1 + \dots + h_n dz_n \rightarrow h_n)$
 $H_0(\Omega^*, X) \cong \mathcal{B}$.

Proof. It follows immediately from the dual version of Corollary 17.12 of [1]. $\square$

Multiplication with c gives us a map of complexes

$$\cdot c: (\Omega^*, X) \rightarrow (\Omega^*, X).$$

We denote by $\mathcal{B}$ the mapping cone associated to this map, see [1]. We have

$$\mathcal{B}: 0 \leftarrow \mathcal{O} \oplus 0 \xleftarrow{\beta} \Omega^1 \oplus \mathcal{O} \xleftarrow{\beta} \dots \xleftarrow{\beta} \Omega^n \oplus \Omega^{n-1} \xleftarrow{\beta} 0 \oplus \Omega^n \xleftarrow{\beta} 0$$

where

$$\beta(\omega, \eta) := \begin{pmatrix} X & (-1)^j c \\ 0 & X \end{pmatrix} \begin{pmatrix} \omega \\ \eta \end{pmatrix} := \begin{pmatrix} X(\omega) + (-1)^j c\eta \\ X(\eta) \end{pmatrix},$$

and j is the degree of $(\omega, \eta) \in \Omega^j \oplus \Omega^{j-1}$.

$$H_0(\mathcal{B}) \cong \frac{\mathcal{B}}{(c)}.$$

Proof. We have an exact sequence of complexes

$$\begin{array}{ccccccc}
& 0 & & 0 & & 0 & \\
& \downarrow & & \downarrow & & \downarrow & \\
0 & \leftarrow \mathcal{R}^0 & \xleftarrow{X} & \mathcal{R}^1 & \xleftarrow{X} & \dots & \xleftarrow{X} \mathcal{R}^n \leftarrow 0 \\
& \downarrow & & \downarrow & & \downarrow & \\
0 & \leftarrow \mathcal{O} & \xleftarrow{X} & \Omega^1 & \xleftarrow{X} & \dots & \xleftarrow{X} \Omega^n \leftarrow 0 \\
& \downarrow & & \downarrow & & \downarrow & \\
0 & \leftarrow \mathcal{O}_{V,0} & \xleftarrow{X} & \Omega_{V,0}^1 & \xleftarrow{X} & \dots & \xleftarrow{X} \Omega_{V,0}^n \leftarrow 0 \\
& \downarrow & & \downarrow & & \downarrow & \\
& 0 & & 0 & & 0 &
\end{array}$$

where $\mathcal{R}^k := f\Omega^k + df \wedge \Omega^{k-1}$. From Lemma 1 one gets the last part of the long exact sequence in homology

$$\begin{aligned}
0 \rightarrow H_2(\Omega_{V,0}^*, X) \xrightarrow{\delta_2} H_1(\mathcal{R}, X) \xrightarrow{i_1} H_1(\Omega^*, X) \xrightarrow{p_1} H_1(\Omega_{V,0}^*, X) \xrightarrow{\delta_1} \\
\xrightarrow{\delta_1} H_0(\mathcal{R}, X) \xrightarrow{i_0} \mathcal{B} \xrightarrow{p_0} \frac{\mathcal{B}}{(f)} \rightarrow 0,
\end{aligned}$$

where δ_1 and δ_2 are the connecting homomorphisms. This gives us

$$H_1(\Omega_{V,0}^*, X) \cong \ker \delta_1 \oplus \text{im} \delta_1.$$

One has $\text{im} \delta_1 = \ker i_0$, $\mathcal{R}^0 = f\mathcal{O}$ and $\mathcal{R}^1 = f\Omega^1 + df\mathcal{O}$. For

$$\omega = f(h_1 dz_1 + \dots + h_n dz_n) + h df$$

we get

$$X(\omega) = f(X_1 h_1 + \dots + X_n h_n) + h c f$$

and therefore

$$\ker i_0 = \frac{\{gf \in \mathcal{O} : gf \in I_n\}}{fI_n + \mathcal{O}(cf)}.$$

We have set $I_n := \mathcal{O}(X_1, \dots, X_n)$ here. On the other hand multiplication gives an isomorphism

$$\cdot f : \frac{\text{ann}_{\mathcal{B}}(f)}{(c)} \rightarrow \frac{\{gf \in \mathcal{O} : gf \in I_n\}}{fI_n + \mathcal{O}(cf)}$$

and this proves

$$\text{im} \delta_1 \cong \frac{\text{ann}_{\mathcal{B}}(f)}{(c)}.$$

Consider $\ker \delta_1$ now. Then we have

$$\ker \delta_1 = \text{imp}_1 \cong \frac{H_1(\Omega^*, X)}{\ker p_1} = \frac{H_1(\Omega^*, X)}{\text{im} i_1}.$$

We have an isomorphism

$$\phi : H_1(\Omega^*, X) \rightarrow \text{ann}_{\mathcal{B}'}(X_n)$$

given as $\phi(h_1 dz_1 + \cdots + h_n dz_n) = h_n$ with $X_1 h_1 + \cdots + X_n h_n = 0$. To prove

$$\ker \delta_1 \cong \frac{\text{ann}_{\mathcal{B}'}(X_n)}{\text{ann}_{\mathcal{B}'}(X_n) \cap \mathcal{B}'(f, \partial f / \partial z_n)}$$

we have to show

$$\phi \circ i_1(H_1(\mathcal{R}, X)) = \text{ann}_{\mathcal{B}'}(X_n) \cap \mathcal{B}'(f, \partial f / \partial z_n).$$

" $\subseteq$ ": Let $\omega \in H_1(\mathcal{R}, X)$, say $\omega = f(h_1 dz_1 + \cdots + h_n dz_n) + h df$ with $X(\omega) = 0$ and so we have $f(X_1 h_1 + \cdots + X_n h_n) + h c f = 0$. ω is mapped to $f h_n + h \partial f / \partial z_n$. Modulo I_{n-1} we have

$$(f h_n + h \partial f / \partial z_n) X_n = f h_n X_n + h c f = 0$$

and this shows the inclusion. Here we have set $I_{n-1} := \mathcal{O}(X_1, \dots, X_{n-1})$.

" $\supseteq$ ": Let $h \in \text{ann}_{\mathcal{B}'}(X_n) \cap \mathcal{B}'(f, \partial f / \partial z_n)$ be a representative, say

$$h X_n = g_1 X_1 + \cdots + g_{n-1} X_{n-1}$$

and $h = s_1 f + s_2 \partial f / \partial z_n$ with $s_1, s_2 \in \mathcal{O}$. It follows that

$$\begin{aligned} s_1 f X_n + s_2 \partial f / \partial z_n X_n &= f(s_1 X_n + s_2 c) - s_2 (\partial f / \partial z_1 X_1 + \cdots + \partial f / \partial z_{n-1} X_{n-1}) \\ &= g_1 X_1 + \cdots + g_{n-1} X_{n-1} \end{aligned}$$

and therefore $f(s_1 X_n + s_2 c)$ is contained in I_{n-1} . This means that $s_1 X_n + s_2 c$ is contained in I_{n-1} , because f is not a zero divisor in $\mathcal{B}'$. Let

$$s_1 X_n + s_2 c = l_1 X_1 + \cdots + l_{n-1} X_{n-1}.$$

Then one has

$$g_1 X_1 + \cdots + g_{n-1} X_{n-1} = f l_1 X_1 + \cdots + f l_{n-1} X_{n-1} - s_2 (\partial f / \partial z_1 X_1 + \cdots + \partial f / \partial z_{n-1} X_{n-1}).$$

Set

$$\omega := -(f l_1 - s_2 \partial f / \partial z_1) dz_1 - \cdots - (f l_{n-1} - s_2 \partial f / \partial z_{n-1}) dz_{n-1} + h dz_n.$$

It follows

$$\omega = f(-l_1 dz_1 - \cdots - l_{n-1} dz_{n-1} + s_1 dz_n) + s_2 \cdot df \in \mathcal{R}^1$$

and

$$\begin{aligned} X(\omega) &= f(-X_1 l_1 - \cdots - X_{n-1} l_{n-1} + s_1 X_n) + s_2 c f \\ &= f(-s_1 X_n - s_2 c + s_1 X_n) + s_2 c f \\ &= 0 \end{aligned}$$

Therefore $w \in H_1(\mathcal{R}, X)$ with $\phi \circ i_1(\omega) = h$. □

2.1 Proof of Theorem 1

The double complex $\mathcal{G}$ We use the construction as in [2]. Consider the double complex $\mathcal{G}$:

Numeration: $\mathcal{G}_{i,0} = \Omega^i$, $\mathcal{G}_{i,1} = \Omega^i \oplus \Omega^{i-1} = \mathcal{G}_{i+k,1+k}$ for $k \geq 1$ and $\mathcal{G}_{i,j} = 0$ for $i < 0$ or $j < 0$. The mappings are $\bar{\alpha}(\omega, \eta) := \pm f\omega + df \wedge \eta$,

$$\alpha(\omega, \eta) := \begin{pmatrix} df & 0 \\ \pm f & df \end{pmatrix} \begin{pmatrix} \omega \\ \eta \end{pmatrix} := \begin{pmatrix} df \wedge \omega \\ \pm f \omega + df \wedge \eta \end{pmatrix}$$

and

$$\beta(\omega, \eta) := \begin{pmatrix} X & \pm c \\ 0 & X \end{pmatrix} \begin{pmatrix} \omega \\ \eta \end{pmatrix} := \begin{pmatrix} X(\omega) \pm c\eta \\ X(\eta) \end{pmatrix}.$$

In [2] it is shown that

$$H_i(\mathrm{tot}\mathcal{G}) \cong H_i(\Omega_{V,0}^*, X) \text{ for } 0 \leq i \leq n$$

and

$$\dim_{\mathbb{C}} H_i(\text{tot}\mathcal{G}) = \lambda \text{ for } i > n,$$

where $\text{tot}\mathcal{G}$ denotes the total complex of $\mathcal{G}$ and λ is defined as in Theorem 1. The argument there does not use at this stage that $(X_1, \dots, X_n)$ is a regular sequence. The homology spaces of the total complex are also obtained by analyzing the spectral sequence (E, d) of $\mathcal{G}$ obtained by taking first the α -homology and then the β -homology.

The spectral sequence (E, d) Let us analyze (E, d) . We get the following 2E -terms by Lemma 1 and Lemma 2:

$$\begin{aligned}
{}^2E_{0,0} &\cong \operatorname{coker} \left(\frac{\mathcal{B}}{(c)} \xrightarrow{\cdot f} \mathcal{B} \right) \cong \frac{\mathcal{B}}{(f)} \\
{}^2E_{0,1} &\cong \ker \left(\frac{\mathcal{B}}{(c)} \xrightarrow{\cdot f} \mathcal{B} \right) \cong \frac{\operatorname{ann}_{\mathcal{B}}(f)}{(c)} \\
{}^2E_{1,0} &\cong \operatorname{coker} \left(H_1(\mathcal{B}) \xrightarrow{\bar{\alpha}} H_1(\Omega^*, X) \right) \\
{}^2E_{1,1} &\cong \operatorname{homology} \left(\frac{\mathcal{B}}{(c)} \xrightarrow{\alpha} H_1(\mathcal{B}) \xrightarrow{\bar{\alpha}} H_1(\Omega^*, X) \right) \\
{}^2E_{l+1,l} &\cong \operatorname{coker} \left(H_1(\mathcal{B}) \xrightarrow{\alpha} H_2(\mathcal{B}) \right) \\
{}^2E_{l,l+1} &\cong \ker \left(\frac{\mathcal{B}}{(c)} \xrightarrow{\alpha} H_1(\mathcal{B}) \right) \\
{}^2E_{l+1,l+1} &\cong \operatorname{homology} \left(\frac{\mathcal{B}}{(c)} \xrightarrow{\alpha} H_1(\mathcal{B}) \xrightarrow{\alpha} H_2(\mathcal{B}) \right)
\end{aligned}$$

where $l \geq 1$ and all other 2E -terms are trivial. The differential on 2E is now given as

$$d_2: {}^2E_{q-1,p+2} \rightarrow {}^2E_{q,p}.$$

Therefore the spectral sequence degenerates at the second level and we are able to compute the total homology of the double complex. First note that

$$\ker \left(H_1(\mathcal{B}) \xrightarrow{\bar{\alpha}} H_1(\Omega^*, X) \right) = \ker \left(H_1(\mathcal{B}) \xrightarrow{\alpha} H_2(\mathcal{B}) \right)$$

by the isomorphisms given in Lemma 1 and Lemma 2. Therefore we get

$$\dim_{\mathbb{C}} H_2(\operatorname{tot} \mathcal{G}) = \dim_{\mathbb{C}} H_i(\operatorname{tot} \mathcal{G}) = \lambda$$

for $i \geq 2$ even. Similarly we get

$$\dim_{\mathbb{C}} H_i(\operatorname{tot} \mathcal{G}) = \lambda$$

for $i \geq 3$ odd. Comparing this with the result on $H_i(\operatorname{tot} \mathcal{G})$ in [2] as stated above and using Lemma 3 for h_1^* Theorem 1 follows.

Corollary 1. *The following algebras are finite dimensional and for n even*

$$\begin{aligned}
\operatorname{ind}_{V,0}(X) &= \dim_{\mathbb{C}} \frac{\mathcal{B}}{(f)} - \dim_{\mathbb{C}} \frac{\operatorname{ann}_{\mathcal{B}}(f)}{(c)} - \dim_{\mathbb{C}} \frac{\operatorname{ann}_{\mathcal{B}'}(X_n)}{\operatorname{ann}_{\mathcal{B}'}(X_n) \cap \mathcal{B}'(f, \partial f / \partial z_n)} \\
&\quad + \dim_{\mathbb{C}} \frac{\mathcal{A}}{(c)} - \dim_{\mathbb{C}} \mathcal{A}
\end{aligned}$$

and for n odd

$$\operatorname{ind}_{V,0}(X) = \dim_{\mathbb{C}} \frac{\mathcal{B}}{(f)} - \dim_{\mathbb{C}} \frac{\operatorname{ann}_{\mathcal{B}}(f)}{(c)} - \dim_{\mathbb{C}} \frac{\operatorname{ann}_{\mathcal{B}'}(X_n)}{\operatorname{ann}_{\mathcal{B}'}(X_n) \cap \mathcal{B}'(f, \partial f / \partial z_n)} + \dim_{\mathbb{C}} \frac{\mathcal{A}}{(f)}.$$

Proof. First note that $\Omega_{V,0}^n \cong \frac{\mathcal{A}}{(f)}$ and so we get

$$\begin{aligned} h_{n-1} &= h_{n-1}^* + \dim_{\mathbb{C}} X(\Omega_{V,0}^n) \\ &= h_{n-1}^* + \dim_{\mathbb{C}} \frac{\mathcal{A}}{(f)} - h_n^*. \end{aligned}$$

Now it follows

$$\sum_{\nu=0}^{n-1} (-1)^\nu h_i = \sum_{\nu=0}^n (-1)^\nu h_i^* + (-1)^{n-1} \dim_{\mathbb{C}} \frac{\mathcal{A}}{(f)}$$

and we get the claim for n odd and for n even

$$\text{ind}_{V,0}(X) = h_0^* - h_1^* - \dim_{\mathbb{C}} \frac{\mathcal{A}}{(f)} + \dim_{\mathbb{C}} \frac{\text{ann}_{\mathcal{A}}(f)}{(c)}.$$

Since we have an exact sequence

$$0 \rightarrow \text{ann}_{\mathcal{A}}(f) \rightarrow \mathcal{A} \xrightarrow{\cdot f} \mathcal{A} \rightarrow \frac{\mathcal{A}}{(f)} \rightarrow 0$$

we get

$$\dim_{\mathbb{C}} \frac{\text{ann}_{\mathcal{A}}(f)}{(c)} - \dim_{\mathbb{C}} \frac{\mathcal{A}}{(f)} = \dim_{\mathbb{C}} \frac{\mathcal{A}}{(c)} - \dim_{\mathbb{C}} \mathcal{A}$$

and the corollary follows. $\square$

Now if $(X_1, \dots, X_n)$ is a regular sequence our formula reduces to the formula of Gomez-Mont in the following way. We find that

$$\lambda = \dim_{\mathbb{C}} \frac{\text{ann}_{\mathcal{B}}(f)}{(c)} = \dim_{\mathbb{C}} \frac{\text{ann}_{\mathcal{B}}(c)}{(f)}$$

and we get for even n

$$\begin{aligned} \text{ind}_{V,0}(X) &= \dim_{\mathbb{C}} \frac{\mathcal{B}}{(f)} - \dim_{\mathbb{C}} \frac{\text{ann}_{\mathcal{A}}(f)}{(c)} + \dim_{\mathbb{C}} \frac{\mathcal{A}}{(c)} - \dim_{\mathbb{C}} \mathcal{A}(c) \\ &= \dim_{\mathbb{C}} \frac{\mathcal{B}}{(f)} - \dim_{\mathbb{C}} \text{ann}_{\mathcal{B}}(f) \\ &= \dim_{\mathbb{C}} \frac{\mathcal{B}}{(f)} - \dim_{\mathbb{C}} \frac{\mathcal{A}}{(f)}. \end{aligned}$$

For n odd we get

$$\begin{aligned} \text{ind}_{V,0}(X) &= \dim_{\mathbb{C}} \frac{\mathcal{B}}{(f)} - \dim_{\mathbb{C}} \frac{\text{ann}_{\mathcal{A}}(f)}{(c)} + \dim_{\mathbb{C}} \frac{\mathcal{A}}{(f)} \\ &= \dim_{\mathbb{C}} \frac{\mathcal{B}}{(f)} + \dim_{\mathbb{C}} \mathcal{A}(c) \\ &= \dim_{\mathbb{C}} \frac{\mathcal{B}}{(f)} - \dim_{\mathbb{C}} \frac{\mathcal{A}}{(c)} + \dim_{\mathbb{C}} \mathcal{A}. \end{aligned}$$

Note that the formula of Gomez-Mont can also be written in this form.

3 Relations to residues and examples

The index of the vector field X can also be computed as a residue, see [5]: Let $\delta_1, \dots, \delta_{n-1} \in \mathbb{R}_{>0}$ be chosen small enough and the real hypersurfaces $\{|X_i| = \delta_i\}$, $i = 1, \dots, n-1$, in general position. Further let the real $(n-1)$ -cycle

$$\Sigma := \{f = 0, |X_i| = \delta_i, i = 1, \dots, n-1\}$$

oriented so that $d(\arg X_1) \wedge \dots \wedge d(\arg X_{n-1}) > 0$. One defines the holomorphic function $\hat{c}$ to be the coefficient of t^{n-1} in the formal power series expansion of

$$\frac{\det(\mathbb{I} - \frac{ti}{2\pi} DX)}{\det(1 - \frac{ti}{2\pi} c)}.$$

Then one has

$$\text{ind}_{V,0}(X) = \int_{\Sigma} \frac{\hat{c} dz_1 \wedge \dots \wedge dz_{n-1}}{X_1 \dots X_{n-1}}.$$

DX is the Jacobi matrix of X here. Let us compute $\hat{c}$ explicitly. Denote by $\sigma_i(DX)$ the coefficients of the characteristic polynomial of DX , i.e.

$$\det(DX - tE) = \sum_{i=0}^n (-1)^i \sigma_{n-i}(DX) t^i.$$

Set $g(t) := \det(E - \frac{ti}{2\pi} DX)$ and $h(t) := 1/(1 - \frac{ti}{2\pi} c)$. We have

$$g^{(k)}(0) = \left(\frac{-i}{2\pi}\right)^k k! \sigma_k(DX)$$

and

$$h^{(l)}(0) = l! \left(\frac{ic}{2\pi}\right)^l.$$

Then we get

$$\begin{aligned} \hat{c} &= \frac{(gh)^{n-1}(0)}{(n-1)!} \\ &= \frac{1}{(n-1)!} \sum_{k=0}^{n-1} \binom{n-1}{k} g^{(k)}(0) h^{(n-k-1)}(0) \\ &= \left(\frac{1}{2\pi i}\right)^{n-1} \sum_{k=0}^{n-1} (-1)^{n-k-1} c^{n-k-1} \sigma_k(DX) \\ &= \left(\frac{1}{2\pi i}\right)^{n-1} \sum_{k=0}^{n-1} (-1)^k c^k \sigma_{n-k-1}(DX). \end{aligned}$$

Using the formula in [4] expressing integrals over cycles in V as Grothendieck residues in $\mathbb{C}^n$ we obtain

$$\text{ind}_{V,0}(X) = \text{res}_{\mathbb{C}^n,0} \left[\frac{\frac{\partial f}{\partial z_n} \sum_{k=0}^{n-1} (-1)^k c^k \sigma_{n-k-1}(DX)}{X_1 \dots X_{n-1} f} \right].$$

If moreover $(X_1, \dots, X_n)$ is a regular $\mathcal{O}$ -sequence we find that there is a $k \in \mathbb{N}$ with $f^k \in I_n$. This means that there is a $\beta \in \mathcal{O}$ with $f^k = \beta X_n \bmod(I_{n-1})$ and therefore we have for any $h \in \mathcal{O}$

$$\begin{aligned} \operatorname{res}_{\mathbb{C}^n, 0} \begin{bmatrix} hc \\ X_1 \dots X_n \end{bmatrix} &= \operatorname{res}_{\mathbb{C}^n, 0} \begin{bmatrix} hc\beta \\ X_1 \dots X_{n-1} f^k \end{bmatrix} \\ \operatorname{res}_{\mathbb{C}^n, 0} \begin{bmatrix} h \frac{\partial f}{\partial z_n} \\ X_1 \dots X_{n-1} f \end{bmatrix} &= \operatorname{res}_{\mathbb{C}^n, 0} \begin{bmatrix} h \frac{\partial f}{\partial z_n} f^k \\ X_1 \dots X_{n-1} f^{k+1} \end{bmatrix} \\ &= \operatorname{res}_{\mathbb{C}^n, 0} \begin{bmatrix} h\beta c f \\ X_1 \dots X_{n-1} f^{k+1} \end{bmatrix} \end{aligned}$$

which means that

$$\operatorname{res}_{\mathbb{C}^n, 0} \begin{bmatrix} hc \\ X_1 \dots X_n \end{bmatrix} = \operatorname{res}_{\mathbb{C}^n, 0} \begin{bmatrix} h \frac{\partial f}{\partial z_n} \\ X_1 \dots X_{n-1} f \end{bmatrix}.$$

We have obtained

Proposition 1. (i) $\operatorname{ind}_{V,0}(X) = \operatorname{res}_{\mathbb{C}^n, 0} \left[\frac{\frac{\partial f}{\partial z_n} \sum_{k=0}^{n-1} (-1)^k c^k \sigma_{n-k-1}(DX)}{X_1 \dots X_{n-1} f} \right]$.
(ii) If $(X_1, \dots, X_n)$ is a regular sequence

$$\operatorname{ind}_{V,0}(X) = \operatorname{ind}_{\mathbb{C}^n, 0}(X) - \operatorname{res}_{\mathbb{C}^n, 0} \begin{bmatrix} \det(DX - c\mathbb{I}) \\ X_1 \dots X_n \end{bmatrix}.$$

We remark that the residues are very easy to compute: If $g_1, \dots, g_n$ is a regular sequence and $h \in \mathcal{O}$ we can find integers $k_1, \dots, k_n$ with $(x_1^{k_1}, \dots, x_n^{k_n}) \subset (g_1, \dots, g_n)$ and therefore a matrix A with

$$\begin{pmatrix} x_1^{k_1} \\ \vdots \\ x_n^{k_n} \end{pmatrix} = A \begin{pmatrix} g_1 \\ \vdots \\ g_n \end{pmatrix}.$$

If d is the coefficient of $x_1^{k_1-1} \dots x_n^{k_n-1}$ in the power series expansion of $h \det A$ then $\operatorname{res}_{\mathbb{C}^n, 0} \begin{bmatrix} h \\ g_1 \dots g_n \end{bmatrix} = d$.

Let us consider an example. $D_k: f = x^2 y + y^{k-1}$, $k \geq 4$.

$$X := \frac{k-2}{2(k-1)} x^{m+1} \frac{\partial}{\partial x} + \frac{1}{k-1} x^m y \frac{\partial}{\partial y}, \quad m \geq 2.$$

We have $c := x^m$ here. The index can be computed easily with Proposition 1 and we obtain $\operatorname{ind}_{V,0}(X) = (m-1)(k-1)$. We want to verify that Corollary 1 gives the same value. Using residues again we compute

$$\dim_{\mathbb{C}} \frac{\mathcal{O}}{(X_1, f)} = (k-1)(m+1).$$

Consider the monomial $x^m y$ in $\frac{\mathcal{O}}{(x^{m+1}, x^2 y + y^{k-1})}$. We find that the monomials $x^m y, \dots, x^m y^{k-2}$ are linearly independent in this algebra and therefore

$$\dim_{\mathbb{C}} \frac{\mathcal{B}}{(f)} = \dim_{\mathbb{C}} \frac{\mathcal{O}}{(X_1, f)} - k + 2 = m(k-1) + 1.$$

Now we claim

$$\{g \in \mathcal{O} : gf \in (X_1, X_2)\} = \mathcal{O}(x^m).$$

Let

$$gf = c_1 x^{m+1} + c_2 x^m y.$$

Then

$$g(x^2 y + y^{k-1}) - c_2 x^m y \in \mathcal{O}(x^{m+1})$$

and therefore

$$g(x^2 + y^{k-2}) \in \mathcal{O}(x^m)$$

since y is not a zero divisor in $\frac{\mathcal{O}}{(x^{m+1})}$. This means $g \in \mathcal{O}(x^m)$ since $x^2 + y^{k-2}$ is not a zero divisor in $\frac{\mathcal{O}}{(x^m)}$. It follows that

$$\dim_{\mathbb{C}} \frac{\text{ann}_{\mathcal{B}}(f)}{(c)} = 0.$$

We also find immediately $\dim_{\mathbb{C}} \mathcal{A} = k$ and $\dim_{\mathbb{C}} \frac{\mathcal{A}}{(c)} = k$. Since $\text{ann}_{\mathcal{B}'}(X_2) = \mathcal{B}'(x)$ we find

$$\frac{\text{ann}_{\mathcal{B}'}(X_2)}{\text{ann}_{\mathcal{B}'}(X_2) \cap \mathcal{B}'(f, \partial f / \partial y)} = \frac{\mathcal{B}'(x)}{\mathcal{B}'(x) \cap \mathcal{B}'(f, \partial f / \partial y)} \cong \mathcal{A}'(x)$$

with $\mathcal{A}' = \frac{\mathcal{O}}{(f, \partial f / \partial y)}$. A basis of $\mathcal{A}'$ is given by $1, x, xy, \dots, xy^{k-2}, y, \dots, y^{k-2}$ and therefore $\dim_{\mathbb{C}} \mathcal{A}'(x) = k$ since $y^{k-2} \sim x^2$ in $\mathcal{A}'$. Now Corollary 1 gives in fact

$$\text{ind}_{V,0}(X) = (m-1)(k-1).$$

References

- [1] EISENBUD, D.: Commutative Algebra. With a View Toward Algebraic Geometry, Graduate Texts in Mathematics 150, Springer-Verlag, New York, 1995
- [2] GOMEZ-MONT, X.: An algebraic formula for the index of a vector field on a hypersurface with an isolated singularity, J. of Alg. Geom. **7**, (1998) 731-752
- [3] GOMEZ-MONT, X.: SEADE, J., and VERJOVSKY, A.: The index of a holomorphic flow with an isolated singularity, Math. Ann. **291**, (1991) 737-751
- [4] KLEHN, O.: Local residues of holomorphic 1-forms on an isolated surface singularity, manuscripta math. **109**, (2002) 93-108
- [5] LEHMANN, D., SOARES, M. and SUWA, T.: On the index of a holomorphic vector field tangent to a singular variety, Bol. Soc. Bras. Mat. **26**, (1995) 183-199